

Références bibliographiques

- **Barrington SF, Johnson PWM. 18F-FDG PET/CT in lymphoma : has imaging-directed personalized medicine become a reality ?** J Nucl Med 2017 ; 58 : 1539-44.
- **Cheson BD, Fisher RI, Barrington SF et al. Recommendations for initial evaluation, staging, and response assessment of Hodgkin and non-Hodgkin lymphoma : the Lugano classification.** J Clin Oncol 2014 ; 32 : 3059-68.
- **Barrington SF, Mikhaeel NG, Kostakoglu L et al. Role of imaging in the staging and response assessment of lymphoma : consensus of the International Conference on Malignant Lymphomas imaging working group.** J Clin Oncol 2014 ; 32 : 3048-58.
- **Haute Autorité de Santé. Évaluation et état des lieux de la tomographie par émission de positons couplée à la tomodensitométrie (TEP-TDM).** Mai 2005.
https://www.has-sante.fr/portail/upload/docs/application/pdf/TEP_TDM_rap.pdf
- **Boellaard R, Delgado-Bolton R, Oyen WJG et al. FDG PET/CT : EANM procedure guidelines for tumor imaging : version 2.0.** Eur J Nucl Med Mol Imaging 2015 ; 42 : 328-54.
- **Valls L, Badve C, Avril S et al. FDG-PET imaging in hematological malignancies.** Blood Rev 2016 ; 30 : 317-31.
- **Meignan M, Gallamini A, Meignan M, Gallamini A, Haioun C. Report on the first international workshop on interim-PET-scan in lymphoma.** Leuk Lymphoma 2009 ; 50 : 1257-60.
- **Luminari S, Biasoli I, Arcaini L et al. The use of FDG-PET in the initial staging of 142 patients with follicular lymphoma : a retrospective study from the FOLL05 randomized trial of the Fondazione Italiana Linfomi.** Ann Oncol 2013 ; 24 : 2108-12.
- **Younes A, Hilden P, Coiffier B et al. International working group consensus response evaluation criteria in lymphoma (RECIL 2017).** Ann Oncol 2017 ; 28 : 1436-47.
- **Trotman J, Luminari S, Boussetta S et al. Prognostic value of PET-CT after first-line therapy in patients with follicular lymphoma : a pooled analysis of central scan review in three multicentre studies.** Lancet Haematol 2014 ; 1 : e17-27.
- **Dreyling M, Ghielmini M, Rule S et al. Newly diagnosed and relapsed follicular lymphoma : ESMO clinical practice guidelines for diagnosis, treatment and follow-up.** Ann Oncol 2016 ; 27 (suppl 5) : v83-90.
- **Raemaekers JM, André MP, Federico M et al. Omitting radiotherapy in early positron emission tomography-negative stage I/II Hodgkin lymphoma is associated with an increased risk of early relapse : clinical results of the preplanned interim analysis of the randomized EORTC/LYSA/FIL H10 trial.** J Clin Oncol 2014 ; 32 : 1188-94.
- **Radford J, Illidge T, Counsell N. et al. Results of a trial of PET-directed therapy for early-stage Hodgkin's lymphoma.** N Engl J Med 2015 ; 372 : 1598-607.
- **André MPE, Girinsky T, Federico M et al. Early positron emission tomography response-adapted treatment in stage I and II Hodgkin lymphoma : final results of the randomized EORTC/LYSA/FIL H10 trial.** J Clin Oncol 2017 ; 35 : 1786-94.
- **Herztberg M, Gandhi MK, Trotman J et al. Early treatment intensification with R-ICE and 90Y-ibritumomab tiuxetan (Zevalin)-BEAM stem cell transplantation in patients with high-risk diffuse large B-cell lymphoma patients and positive interim PET after 4 cycles of R-CHOP-14.** Haematologica 2017 ; 102 : 356-63.
- **Casasnovas RO, Ysebaert L, Thieblemont C et al. FDG-PET-driven consolidation strategy in diffuse large B-cell lymphoma : final results of a randomized phase 2 study.** Blood 2017 ; 130 : 1315-26.
- **Kanoun S, Rossi C, Berriolo-Riedinger A et al. Baseline metabolic tumour volume is an independent prognostic factor in Hodgkin lymphoma.** Eur J Nucl Med Mol Imaging 2014 ; 41 : 1735-43.
- **Cottreau AS, Becker S, Broussais F et al. Prognostic value of baseline total metabolic tumor volume (TMTV0) measured on FDG-PET/CT in patients with peripheral T-cell lymphoma (PTCL).** Ann Oncol 2016 ; 27 : 719-24.